

IMPORTANT MEDICAL INFO



**THIS PATIENT LACKS PARATHYROID HORMONE
AND IS AT RISK FOR HYPOCALCEMIC AND
HYPERCALCEMIC CRISES.**

In case of serious illness, nausea, vomiting and/or cramps, check serum calcium and kidney function.

If hypocalcemia and tetany, administer i.v. 2.5-5 mmol calcium chloride or gluconate (~ 100-200 mg elemental calcium) in 100 mL saline or glucose solution over 10 min. Repeat if needed.

If hypercalcemia > 3.5 mmol/L (14.0 mg/dL), administer saline intravenously and admit the patient immediately to hospital.

Name/ Ime

Date of birth/ Datum rojstva



European Society
of **Endocrinology**

POMEMBNO ZDRAVNIŠKO OBVESTILO



BOLNIKU PRIMANJUJE PARATHORMONA, ZATO JE OGROŽEN ZA NASTANEK HIPOKALCEMIČNE IN HIPERKALCEMIČNE KRIZE.

V primeru resne bolezni, slabosti, bruhanja in/ali krčev pri bolniku preverite serumski kalcij in delovanje ledvic.

V primeru hipokalcemije in tetanije bolniku aplicirajte 2,5–5 mmol kalcijevega glukonata ali klorida (pribl. 100–200 mg elementarnega kalcija) v 100 ml fiziološke raztopine ali raztopine 5% glukoze v 10 minutah. Po potrebi ponovite.

V primeru hiperkalcemije >3,5 mmol/l bolniku nastavite intravensko infuzijo fiziološke raztopine in ga takoj hospitalizirajte.



ZDRUŽENJE
ENDOKRINOLOGOV
SLOVENIJE